

GLYCERIN SUPPOSITORIES 2g CHILDREN'S SIZE

Read all the carton before use.

What the medicine is for

Glycerin Suppositories relieve occasional constipation by helping to stimulate bowel movement.

Before you use Glycerin Suppositories

X Do not use the medicine if your child has....

- An **allergy** to any of the ingredients.
- Do not use for adults or infants under 1 year old.

✓ How to use the medicine

- Suppositories must not be taken by mouth.
- Wash your hands before and after use.
- Remove the protective film before use by tearing the strip at the notch or by cutting around the suppository.
- Dip the suppository in water and insert into the rectum.

Children: Insert one suppository as required.

Do not use for adults or infants under 1 year old.

If symptoms persist ask your doctor for advice.

! Not to be swallowed. If anyone accidentally swallows some see a doctor straight away.

Possible side effects

May cause bowel irritation, and occasionally abdominal cramps. If your child has any side effects, stop use and tell your doctor or pharmacist.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

Storing the medicine

Do not store above 25°C. Store in a dry place.

KEEP OUT OF THE SIGHT AND REACH OF CHILDREN.

Use by the date shown on the side of the carton.

Active ingredient per suppository: glycerol 1.4g.

Also contains: gelatin and purified water.

MA holder and manufacturer: Thornton and Ross Ltd., Huddersfield, HD7 5QH, UK.

Text revised: November 2014

PL 00240/6101R

22467509