200

<complex-block><text><text><list-item><list-item><list-item><list-item><list-item><list-item><list-item><list-item><list-item><list-item><text><text><text><text><text><text><text><text><text><text><text><text><text><text><text><text></text></text></text></text></text></text></text></text></text></text></text></text></text></text></text></text></list-item></list-item></list-item></list-item></list-item></list-item></list-item></list-item></list-item></list-item></text></text></complex-block>	12 mm		150 mm		12 mm	
		<ul> <li>Finis medicine is for the short term treatment of acute moderate pain when other painkillers have not worked.</li> <li>You should only take this product for a maximum of 3 days at a time. If you need to take it for longer than 3 days you should see your doctor or pharmacist for advice.</li> <li>This medicine contains codeine which can cause addiction if you take it continuously for more than 3 days. This can give you withdrawal symptoms from the medicine when you stop taking it.</li> <li>If you take ting through this leaflet before you start using this medicine.</li> <li>This medicine is available without prescription, but you still need to use Solpadeine Max Tablets carefully to get the best results from them.</li> <li>Hyou have any questions, or if there is anything you do not understand, ask your pharmacist.</li> <li>In this leaflet</li> <li>Mow to stake Solpadeine Max Tablets</li> <li>How to store Solpadeine Max Tablets</li> <li>How to store Solpadeine Max Tablets.</li> <li>Fousible side effects</li> <li>How to store Solpadeine Max Tablets do</li> <li>Check before you take Solpadeine Max Tablets over 12 years of age for the short-term relief of moderate pain that is not relieved by other painkillers such as paracetamol or ibuprofen alone. They can be used for migraine, headache, dental pain, period pain, strains &amp; sprains, backache, arthritic &amp; rheumatic pain and sciatica. The tablets contain two active ingredients; paracetamol and codeine are both painkillers. Codeine belongs to a group of medicines allel opioid analgesics which act to relieve pain. It can be used on its own or in combination with other painkillers.</li> <li>Check before you take Solpadeine which can cause addiction if you take it continuously for more than 3 days.</li> <li>This medicine contains codeine which can cause adjacent when you stop taking it.</li> <li>If you take a painkiller for headaches for more</li> </ul>	<ul> <li>If you have ever had an allergic reaction to paracetamol, codeine, other opioid painkillers or to any of the other ingredients (listed in Section 6)</li> <li>If you are taking other medicines containing paracetamol or codeine</li> <li>For pain relief in children and adolescents (0-18 years of age) after removal of their tonsils or adenoids due to obstructive sleep apnoea syndrome</li> <li>If you know that you metabolise very rapidly codeine into morphine</li> <li>If you are breastfeeding.</li> <li>Ask your doctor before you take this medicine:</li> <li>If you have liver or kidney disease, including alcoholic liver disease</li> <li>If you have bowel problems including blockage of your bowel</li> <li>If you have bad an operation to remove your gall bladder</li> <li>If you have been told by your doctor that you have an intolerance to some sugars.</li> <li>If you are taking other medicines</li> <li>Talk to your doctor or pharmacist before taking these tablets if you are taking any prescribed medicines; particularly metochoramide or domperidone (for nausea [feeling sick] or vomiting [being sick]); colestyramine (to lower blood cholesterol); monoamine oxidase inhibitors (MAOIs) prescribed for depression or medicines that make you drowsy. If you take blood thinning drugs (anticoagulants e.g. warfarin) and you need to take a pain reliever on a daily basis, talk to your doctor because of the risk of bleeding. But you can still take occasional doses of Solpadeine Max Tablets at the same time as anticoagulants.</li> <li>Warnings and precautions</li> <li>Codeine is transformed to morphine in the liver by an enzyme. Morphine is the substance that produces pain relief. Some people have a variation of this enzyme and this can affect people are more likely to get serious side effects because a very high amount of morphine is produced. If you notice any of the following side effects, you must stop taking this medicine and adolescents after surgery: Codeine should</li></ul>	Ph Code 12804 80 mm	<ul> <li>Pregnancy and breast feeding Talk to your doctor before taking Solpadeine May Tablets wile you are pregnant. Do not take Solpadei Max Tablets wile you are breast-feeding. Codein and morphine passes into breast milk.</li> <li>Driving and using machines This medicine can affect your ability to drive as it may make you sleepy or dizzy. Do not drive while taking this medicine until you know how it affects you. It is an offence to drive if this medicine affects you ability to drive. However, you would not be committing an offence if. • The medicine has been taken to treat a medical dental problem and • You have taken it according to the information provided with the medicine and • I vas not affecting your ability to drive safely. Talk to your doctor or pharmacist if you are not su medicine. Details regarding a new driving offence concernit driving after drugs have been taken in the UK ma be found here: www.gou.uk/drug-driving-law</li> <li>3. How to take Solpadeine Max Tablets Adults: Swallow 2 tablets every 4-6 hours as needed. Do not take more frequently than every 4 hours. Do not take more frequently than every 6 hours. Do not take more than 8 tablets in 24 hours. Do not take more than 8 tablets in 24 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hours. Do not take more frequently than every 6 hou</li></ul>	<ul> <li>Constipation</li> <li>Stop taking this medicine and tell immediately if you experience:         <ul> <li>Severe abdominal pain, nausea a have recently had your gall blad</li> <li>Allergic reactions which may be rash and itching sometimes with mouth or face or shortness of br</li> <li>Skin rash or peeling or mouth ul</li> <li>Breathing problems. These are m have experienced them before v painkillers such as ibuprofen and</li> <li>Unexplained bruising or bleedin</li> <li>Nausea, sudden weight loss, loss yellowing of the eyes and skin.</li> </ul> </li> <li>Reporting of side effects: If you gat talk to your doctor, pharmacist or includes any possible side effects r leaflet. You can also report side eff the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard</li> </ul> <li>By reporting side effects you can h information on the safety of this r</li> <li>How do I know if I am addicted if you take this medicine according instructions on the pack, it is unlik become addicted to the medicine. following apply to you it is import to your doctor:         <ul> <li>You need to take more than the dose.</li> <li>When you stop taking the medic unwell but you feel better if you medicine again.</li> </ul> </li> <li>How to store Solpadeinee Keep out of the reach and sight o Do not use this medicine after the on the pack.</li> <li>Further information Active ingredients Each tablet con 500 mg and Codeine Phosphate He Other ingredients Starch pregelati maize starch, talc, magnesium stee microcrystalline cellulose, croscarm inactose monohydrate, hypromello quinoline yellow (E 104), erythrosi titanium dioxide (E 171). Packs of Solpadeine Max Tablets or soth sides.</li> <li>matheting authorisation hold Pharma Ltd, 32 Vauxhall Bridge Re 2SA, United Kingdom and all enqu sent to this address.</li> <li>The manufactures is Omega Pharm GmbH &amp;</li>

Note: Main body font size 7pt/7.5pt

